JH AITC - ALL COHORTS



Predicting Adverse Events from Sovrinti In-Home ADL Sensing



AREA OF NEED:

The need to identify rising health risks in Aging individuals to enable interventions

before the issue becomes acute.

OPPORTUNITY:

Develop and test, change models that use in-home ADL sensors to identify and predict rising risk of acute conditions in

Aging individuals.

DATA:

From recently completed NIA study: 117 ADRD Caregiver-Care recipient dyads with an average of 12 months of in-home sensors data and coincident monthly ADCS surveys and health conditions information. 800 reported health incidents

TECH APPROACH:


Leverage real time location and device utilization data to identify behavioral anomalies associated with described incidents. Utilize various ML/AI techniques to develop a predictor of acute conditions from sensor data.

PI(s): John Fitch

COHORT: GY1



AI/ML Based ADL Change Detection



Unique Data of Daily Living™



WAVi Ai: Preventing Age-Related Cognitive Decline



Age-related cognitive decline can be **AREA OF NEED:**

mitigated for many patients with early identification that includes interventional

phenotyping.

Create a software product to be added to **OPPORTUNITY:**

> the existing WAVi EEG/ERP platform intended to be used in the setting of age-

related cognitive decline as both a

diagnostic aid and to track interventions.

Analysis will be on Existing WAVi data DATA:

collected in clinic on patients concerned

with or experiencing age-related


declines, including longitudinal outcomes.

Add an Ai component to the standard **TECH APPROACH:**

> EEG/ERP metrics on a heterogenous data set in order to find common feature sets that both identify early dementia and help predict candidate interventions.

PI(s): Francesca Arese, PhD David Oakley, PhD







RetreatVR – A Senior Wellness Tool



AREA OF NEED: Mitigating senior loneliness through Virtual

Reality

OPPORTUNITY: Develop, test, and implement shared

immersive experiences to enhance social

interaction

DATA: Correlation study analyzing the

correlation between the level of

engagement with RetreatVR and the

level of social engagement

TECH APPROACH: Designed and developed a senor-friendly

VR platform that included immersive and

interactive 3D experiences to tap memories and encourage social

interaction

PI(s): Ellie Giles





Figure 1: Eye-gaze-activated menu



Robot + Al-based Facial Expression Analysis to Detect Agitation in Persons with AD/ADRD



AREA OF NEED:

Physical and verbal agitation, found in 80% of nursing home residents with AD/ADRD, can pose a major problem for the resident and their family and professional caregivers. The inability to successfully manage dementia behaviors often results in increased burden on caregivers, use of physical restraints, and/or pharmacological interventions. Use of antipsychotic and psychotropics medications increases the risk of death and of falls and fractures in patients with dementia.

OPPORTUNITY

Integrate an AI-based facial expression analysis software on an autonomously navigating robot to test and validate that it can detect agitation in nursing home residents with dementia. The ultimate goal is to use this information and the robot to deliver non-pharmacological interventions to reduce agitation.

DATA:

Collecting comprehensive raw data of facial expressions of 10 nursing home residents with dementia known to be agitated during three days of simultaneous observations twice per hour for 13 hours per day by the robot and two research assistants.

TECH APPROACH:

Using the ability of the autonomously navigating robot to come to a resident's room in a nursing home as frequently and for as long a time as necessary, and through its video camera capture facial expressions and use the facial expression analysis software to analyze the images and validate that a signal can be seen when a resident is agitated as determined by the research assistants.

PI(s):

Yuval Malinsky, MD


COHORT:

GY1





Implementing the Emergency Department Dementia Algorithm (EDDA) for Dementia Detection in At-Risk ED Patients



AREA OF NEED:

The ED is underutilized as an environment where screening and identification of undiagnosed people living with dementia can occur.

OPPORTUNITY:

Train and validate the EDDA machine learning algorithm as a clinical decision support tool that will screen at-risk ED patients for dementia upon visit.

DATA:


Demographic, diagnostic, laboratory, procedural, and medication data from all ED patients seen from 2013-2020 in the Yale New Haven Hospital System.

TECH APPROACH:

Integrate the EDDA into the EHR to convey real time identification alerts for ED providers, allowing for earlier interventions for undiagnosed persons living with dementia.

PI(s):

Ula Hwang MD, MPH Andrew Taylor MD, MHS





Al Bots and Wearables for Dementia Caregivers to Improve Quality of Life



AREA OF NEED: A previous study by People Power Company dba Care Daily, evaluated a sensor-based in-home support system and found a significant decrease in caregiver anxiety in the active group as compared to the control group1. Dementia caregivers experience higher levels of anxiety than non-caregivers and anxiety is associated with suicidal ideation.

OPPORTUNITY:

This improved RCT study substitutes 2 Apple Watches in place of 12 sensors; substitutes a 5-minute weekly voice survey in place of a 90-minute quarterly survey using Qualtrics; and upgrades

wellness reports using HIPAA-compliant ChatGPT.

Al Bot Features: Alert on falls, Detect wandering, Alert with GPS location, Monitor sleep quality trends, Monitor fitness, Caregiver

blog with wellness tips for caregivers in need

DATA:

Participants: Goal of 40 dyads (each with family Cg and PWD). Currently 93 dyads applied, 38 dyads consented. A 4 Month

Randomized Controlled Trial (RCT) is underway collecting data from 19 of 20 active caregivers who received 2 Apple Watches (for CG

and PWD) and sent 5-minute weekly wellness surveys The control group with 19 of 20 CGs did not receive Apple

Watches but are sent complete 5-minute weekly wellness surveys

TECH APPROACH:

Measures: Self-reported CG anxiety, depression, and well-being checked weekly using voice survey and biometrics captured with

Apple Watches. Procedure/Timeline:

> July 24 through Dec 31, 2023: RCT conducted January 2024: Data analysis and reporting

Gene Wang. Contact: www.caredaily.ai

COHORT:















Balance T to Improve Balance and Reduce Fall Risk



AREA OF NEED: Falls are the leading cause of injury death

> among adults > 65. The Incidence of Falling is increasing over the last 10 years. Most

people fall while turning or walking

Develop, test, and validate the use of the **OPPORTUNITY:**

> Balance-T to improve balance skills and reduce risk for falls of older adults at

home and in the community.

Collecting data through functional DATA:

outcome measures of balance and gait

TECH APPROACH: Implement means to track compliance and motion of both Balance T and body

via the user's cell phone and wearable

technology (i.e. smart watch).

Balance T is custom fit to each user, for height, shoulder-width, and degree of

difficulty (handlebar tilt)

PI(s): Michael Schubert, PT PhD

Yuri Agrawal MD

COHORT: Pre-Frail and Robust Older Adults

Community Dwelling

IMPROVE YOUR BALANCE AND AGILITY



These exercises are designed to improve balance whether you are young, old, injured or healthy



Booklet includes 4 exercises to improve balance



Balance T enables balance training during standing and walking!



Individualized risk evaluation of cognitive decline in a cognitively normal population



AREA OF NEED: Subjective cognitive decline (SCD) is a group at high risk for dementia, but

there is no method to predict dementia outcomes on an individual basis.

There are also no guidelines to prevent conversion from SCD to dementia.

OPPORTUNITY: Develop, test, and validate machine learning

models tailored for individuals with SCD to

predict potential cognitive decline on a

personalized basis, and to pinpoint modifiable risk

factors associated with this deterioration.

DATA: Approximately 600 individuals without dementia

underwent clinical evaluations, which included

cognitive assessments and baseline brain MRI scans,

and were subsequently monitored for 2 years.

TECH APPROACH: Measure the local brain volumes from MRI scans, which will then be integrated with clinical data,

including the presence of modifiable risk factors associated with the onset of dementia. Identify

brain anatomical features and clinical markers linked to cognitive decline over a two-year period.

The final model will be tested on real-world clinical data to assess its relevance to a diverse clinical


population.


PI(s): Kenichi Oishi, MD, PhD

COHORT: GY1



Building models for risk prediction using existing data







Leveraging Conversational AI to Detect Cognitive Impairment and Dementia in the Home



AREA OF NEED: Few validated, scalable approaches to early

detection of AD/ADRD exist, limiting

potential to introduce targeted

pharmacologic and lifestyle interventions

OPPORTUNITY: Develop, test, and validate machine

learning models that use conversational Al to identify and accurately predict the cognitive status of older adults at home


TECH APPROACH: Leverage consumer-friendly voice

assistant technologies (e.g., Amazon, Google) equipped with scientifically validated, clinical-grade cognitive assessments and a personalized digital coach to provide and coordinate support

DATA: Collecting comprehensive cognitive

assessment data, including raw audio data, from 160 patient-caregiver dyads

PI(s): Randall Williams, MD





Visilant: Equitable Access to Eye Care Through Telemedicine and Artificial Intelligence



AREA OF NEED: Lack of access to cataract screening prevents

disadvantaged populations from seeking care, leading to disease progression, functional

limitations, and worse outcomes

OPPORTUNITY: Develop a simple and inexpensive anterior

segment imaging and telemedicine system to allow for remote eye screening facilitated by

non-ophthalmologists

DATA: Project will build a database of at least 2,200

images of eyes with no cataract, immature cataract, and mature cataract and validate a diagnostic Al algorithm in 100 patients against


a gold-standard exam

TECH APPROACH: Visilant system includes a mobile app with Al

and telemedicine capabilities, a proprietary smartphone attachment for ocular imaging, and a patient management dashboard and provider portal. The ML approach uses eye segmentation, classification with a CNN, and domain adaptation technique to mitigate bias.

PI(s): Kunal Parikh, PhD

Nakul Shekhawat, MD, MPH



Proprietary smartphone attachment for simple, high quality anterior segment imaging



Mobile app with AI and telemedicine capabilities



Patient management dashboard and provider portal



Enhancing Slow-Wave Sleep in Older Adults



AREA OF NEED: Growing prevalence of dementia in older adults

highlights a pressing need for early-stage therapies. Specifically, there's a noticeable gap in therapies targeting the underlying pathology of sleep disruptions

in patients experiencing cognitive challenges.

OPPORTUNITY: To take a lab-based technique, which modulates critical

brain oscillations, and translate it into a non-invasive wearable technology for everyday at-home use. This technology would focus on enhancing slow wave sleep

using closed-loop neuromodulation.

DATA: Collecting comprehensive sleep EEG data, from 20

older adults totaling 150 nights of data.

TECH APPROACH: The key lies in precision. By employing sound as the


medium, our closed loop technology is designed to

deliver precise stimulation, aiming to optimize oscillations during deep sleep, thereby enhancing its

restorative effects on cognition.

PI(s): Joshua Blair, MS

Spencer Shumway, MS Youseph Yazdi, PhD







Al Driven Avatar in DevaWorld, a Dementia Friendly Virtual World



AREA OF NEED: While the benefits of a person-centered care

approach for people living with dementia (PLwD) are well documented, few caregivers have the time (staff shortage) or the conversational and relationship skills

to fully engage with PLwD on a daily basis.

OPPORTUNITY: Develop, test, and validate a conversational Al agent

to fully engage with PLwD without adding caregiver burden, nor relying on specialized caregiver skills.

DATA: Video data of PLwD using our Al version of

DevaWorld, along with emotion recognition analyses

and tapping patterns withing the virtual world.

TECH APPROACH: Use a mix of rule-based, statistical-based and prompt

engineering approaches; integrated with speech-to-text (STT), facial and speech emotion recognition.

PI: Algis Leveckis, SM







Artificial Intelligence Algorithm to Improve Palliative Care of Alzheimer's Patients



CENTER FOR POPULATION HEALTH

AREA OF NEED:

Providing adequate palliative care for individuals with AD/ADRD presents numerous challenges such as determining a patient's status on the AD/ADRD disease course, creating need for improved methods of identifying patients who would benefit from timely palliative care (PC) in the community.

OPPORTUNITY:

Develop and validate machine-learning predictive models to identify AD/ADRD patients who are likely to benefit from PC assessment.

DATA:

Electronic health record (EHR) data of patients with AD/ADRD receiving care in the Johns Hopkins Health System over 2017–2021.


TECH APPROACH:

ML prediction algorithm incorporating rich clinical information available in EHR data from academic healthcare system to identify persons with AD/ADRD who would most benefit from PC, and its impact on healthcare utilization outcomes and disparities in PC delivery.

PI(s):

Chintan J. Pandya MD, PhD Jonathan Weiner, DrPH







Quantifying the features and predictors of cognitive fluctuation in Alzheimer's Disease and MCI

AREA OF NEED: Cognitive fluctuations, defined as

spontaneous alterations in attention, arousal, and cognition, have been assessed anecdotally. However, the exact features and causes of cognitive fluctuations are still

unclear.

OPPORTUNITY: Ecological momentary assessments is

crucial to determining the features of cognitive fluctuations by remotely

measuring memory, attention, and other

executive functions.

DATA: Collecting comprehensive cognitive

assessment data, including raw audio data, from 90 patient-caregiver dyads (30

dementia, 30 MCI, 30 Control)

TECH APPROACH: Utilizing a three-dimensional view of


cognition that combines tablet-based cognitive testing and caregiver reports with monitoring of biomarkers with

wearables.

Kishore Kuchibhotla, Ph.D

COHORT: GY1

PI(s):





EZ-Aware: Digital Twin for Wearable-Enabled, Al-Supported Detection of Cognitive Impairment



AREA OF NEED:

"Baby boomers" (55+) make up 30% of the U.S., or 97.5 million people. As this generation ages, the prevalence of Alzheimer's disease (AD) and related dementias (ADRD) in the U.S. is projected to surge to 12.7 million by 2050. Subjective cognitive decline (SCD) is an early ADRD symptom. While 50-80% of those 65+ report decline, fewer than half are confirmed with impairments, highlighting the urgency for accessible and precise MCI screening methods.

OPPORTUNITY:

Develop and test an accessible digital health platform supported by digital twin models for monitoring of cognitive and behavioral functions in older adults' everyday environments.

DATA:

A 6-week study on 30 older adults, collecting comprehensive active and passive daily life data (using an app connected to a wearable) on cognitive and behavioral functions.

TECH APPROACH:

Integrating smart wearables and schedulable cognitive and behavioral assessments at home provides quality of life datasets used for personalized insights on an individuals ongoing cognitive status.

PI(s): Kunal Mankodiya, PhD





Neuroanatomic validation of digital voice



AREA OF NEED: Digital voice recordings offer novel

capabilities with great potential for early detection of cognitive and associated

functional change.

OPPORTUNITY:

For voice technologies to reach wide adoption, one needs to systematically validate them against known markers of disease and identify the settings where they confer benefit over existing modalities.

DATA:

Our access to thousands of voice recordings, MRI, and pathology data from the Framingham Heart Study puts us in a unique position to build and validate voice-based technologies.


TECH APPROACH:

- We are developing interpretable and generative deep learning approaches to assess cognition from digital voice recordings.
- We have recently formed <u>Vakta.ai</u>, a spin-out from Boston University, which is commercializing our technology.

Pl(s): Vijaya B. Kolachalama, PhD

Rhoda Au, PhD

COHORT: GY2



Schematic of our deep learning framework used for assessing cognition on voice recordings.

[1] Karjadi et. al., 2023. Journal of Alzheimer's Disease



transcription

high quality speechto-text conversion with time-stamped words

diarization

categorize audio content by identifying different speakers

translation

multi language support for speech-totext, text-to-text translation

prediction

text summarization, sentiment analysis, assessment of cognitive status



A Novel Insole Solution Used in Daily Life to Identify and Mitigate Falls and Frailty



AREA OF NEED: Need for real-time data providing insight in

how to treat or track fall risk. Falls are associated with immobility, mortality, and decreased independence; 1 of 4 individuals over 65 years old suffers a fall each year.

OPPORTUNITY: Validate a new portable pressure sensing

insole technology, enabling more efficient and effective collection of clinically relevant balance data to predict

and treat falls in the elderly.

DATA: Collect gait and balance data from 50

individuals > 50 years old and compare insole data to gold standard. Conduct focus groups to validate usability of data.

TECH APPROACH: All algorithms will be used to develop fall

risk classification models utilizing insole data and gold standard parameters. The data will be utilized to develop a protocol for remote therapeutic monitoring to assist in the prevention of falls.

PI(s): Linda Denney, PT, PhD, MAppSc (Manip)

Dan Peterson, PhD







Geriatric Functional Assessment System Using Passive Wearable Sensing and Deep Learning



AREA OF NEED:

Clinics do not routinely assess physical function, and survey assessments that are used by some are

subject to recall and measurement bias

OPPORTUNITY:

Develop, test, and validate a wearable device that will accurately assess physical

function of older adults

DATA:

The device is used to capture video images and collect motion data of an individual as they perform a series of physical function assessments while wearing the device

TECH APPROACH:

An Arduino Vision Shield is used to capture images, and an Arduino IMU Shield for collecting motion data

PI(s):

John A. Batsis, MD Xiaohui Liang, PhD







Machine learning to predict post-COVID-19 cognitive decline and dementia



AREA OF NEED: COVID-19 illness may increase the risk for

Alzheimer's Disease. There is a pressing need to develop novel, accessible, sensitive methods of predicting post-COVID-19 cognitive decline and dementia risk in order to allow for early

detection and intervention

Utilize remote, app-based measurement of **OPPORTUNITY:**

> cognition to develop a more accurate, reliable, and accessible method to detect and predict cognitive

dysfunction and dementia risk

Collecting (1) longitudinal app-based cognitive DATA:

> data, (2) validated symptom inventories, and (3) electronic medical record data from 120 older

adults following COVID-19 illness


Applying machine learning to the DANA app's **TECH APPROACH:**

> rich, high-frequency, high-sensitivity data to develop sensitive predictive algorithms

PI(s): Tracy Vannorsdall, PhD

COHORT:







Automatic Assessment of Neuropsychiatric Symptoms Using Non-Intrusive Contactless Ambient Intelligence Technologies



AREA OF NEED: Neuropsychiatric symptoms (NPS) in dementia

patients are often missed due to subjective assessments based on clinician observation and caregiver interviews. Accurate methods for

assessing and monitoring NPS are urgently needed.

OPPORTUNITY: With better detection of not only NPS,

and the more pervasive Mild Behavioral Impairment, there is potential to improve care and optimize the quality of life for

patients at risk of MCI and AD.

DATA: We are recruiting participants with a

history of behavioral changes and installing sensors in their homes to monitor their NPS for up to 90 days.

TECH APPROACH: We will develop a Computer Vision NPS Assessment

(CVNA) system using ambient intelligence to provide

objective and personalized detection of NPS subsyndromes across the preclinical and clinical spectrum of dementia. To quantify NPS, we adopt the Mild Behavioral Impairment Checklist (MBI-C).

PI: Ehsan Adeli, PhD

Co-ls: Christine Gold, PhD; Vankee Lin, PhD, RN





EyeControl AI: Detecting and Managing Delirium to Reduce Cognitive Decline in Older ICU Patients



AREA OF NEED: Delirium in ICU patients, especially older adults, is a major problem

resulting in poor outcomes including increased mortality, longer hospital stays, higher rates of cognitive decline, and increased healthcare costs. Effective communication and reorientation interventions can help reduce

delirium.

OPPORTUNITY: Test the EyeControl communication device and Al

platform to detect delirium using an automated CAM-

ICU and manage delirium through frequent

communication and orientation messages in ICU

patients.

DATA: Demographic, clinical, delirium assessment (CAM-ICU), cognition

(IQCODE, MoCA), and satisfaction data will be collected on 30 older ICU patients using EyeControl at Johns Hopkins Bayview

Medical Center.

TECH APPROACH: The EyeControl wearable eye-tracking device and Al platform

will enable ICU patients to answer orientation questions and receive communication interventions. The automated CAM-ICU will be compared to standard assessments to evaluate

accuracy. Delirium duration, cognitive outcomes, and

satisfaction will be assessed.

PI(s): M. Haroon Burhanullah, MD Paul B Rosenberg, MD





Predicting Fall Risk in Older Adults Using Machine Learning



AREA OF NEED: Early detection of cognitive decline through the

use of interventional phenotyping can support preventive action to preserve brain health and

mitigate fall risk

OPPORTUNITY: Use point of care data to create an algorithm to

predict falls and provide real time feedback to

clinicians

DATA: Balance data with eyes open and closed will be

collected while performing intake vitals in the HSC clinic for older adults on all patients to create the

predictive algorithm.

TECH APPROACH:


Beyond standard supervised ML predictive models, the data collected will be used to generate an unsupervised autoencoder network that learns statistical relationships between demographics, medical comorbidities, balance variables, and falls that can be used for anomaly detection, missing data inference, and improved prediction using the latent

representational learned.

Pl(s): Rita M. Patterson, PhD, Mark Albert, PhD.,

Kathlene Camp, MPT, DPT, Kim Fulda, Dr. P.H.,

Janice Knebl, DO, MBA.





Picasso Intelligence: Improving Mobility for Dementia Alleviation in Older Adults via Al-Powered Affordable Exosuits

AREA OF NEED: Reduction in ph

Reduction in physical activity have detrimental effects on the health of older adults as it initiates a cycle of declining health which increases the risk of Alzheimer's disease and related dementias (ADRD)

OPPORTUNITY:

Leverage lightweight and affordable exosuits in concert with learning-based personalized control to promote physical activities in everyday settings and ultimately decrease the risk of cognitive decline which leads to dementia in older adults.

DATA:

Analysis will be on data collected in human studies in community settings involving 20 older adults aged 65+ over 9 visits.

TECH APPROACH:

Deep reinforcement learning framework with the actor-critic method and high-fidelity musculoskeletal modeling method to automatically train the exosuit controller to generate continuous assistive torque during versatile activities common in community settings

PI(s):


Chien-Ming Huang, PhD Hao Su, PhD, Junxin Li, PhD

1. Reduced Physical Activity Exosuit walking 3. Cognitive Decline intervention ultimately leading to ADRD 3. Increased Strength, 2. Gait learning Fitness control Physical Activity, **Cognitive Function** 3. Joint Degradation & Pain Increased 2. Reduced Strength & Fatigue Fitness











Smart Rep, Smart Automation to Respond to Patient portal messages

AREA OF NEED: GPT is increasingly used in healthcare

messaging. Thoughtful application of GPT in messaging related to ADRD may be beneficial, as care partners are neither routinely identified nor well supported.

OPPORTUNITY: Develop and test "Smart Rep", which

optimizes responses to patient portal messages to both increase recognition of care partners, and support care partners with vetted content and resources.

DATA: Patient portal messages sent from

persons with dementia and their care partners, and qualitative data collected

from interviews


TECH APPROACH: Through prompt engineering, guide GPT

to a) recognize when a care partner has authored a message on a patient's behalf,

and b) draw on vetted ADRD best

practices and care resources

Pl(s): Kelly T. Gleason, PhD, RN





ADPIE: Deep phenotyping of people with AD using portable integrated equipment

AREA OF NEED: Most digital biomarkers do not leverage the

potential complementarity of multiple measures. Also, most approaches are developed in laboratory conditions, illsuited for deployment to clinical spaces.

OPPORTUNITY: Validate the feasibility of portable integrated equipment (PIE) to obtain

multimodal digital biomarkers of AD.

DATA: Collecting eye movement, speech, and


handwriting in PIE and also laboratory conditions, from 120 participants.

TECH APPROACH: A multimodal platform integrating an

eye-tracker, microphone, and digital pad. Multimodal physiological measurements are analyzed after recording to provide digital biomarkers specific of AD.

PI(s): Laureano Moro-Velazquez, PhD

COHORT: GY3



High Precision Equipment







Al Enhanced Scheduling to Enable an Affordable Neighborhood Model for In-Home Caregiving

AREA OF NEED: The number of seniors in need of home care exceeds the workforce of

caregivers by 6:1. Current staffing models are inefficient and unaffordable resulting in a growing gap group of 18 million seniors without the ability stay healthy at home and stressing the nation's healthcare finances, private assets, and employers of family caregivers.

OPPORTUNITY:

The neighborhood model of home care improves efficiency by enabling a caregiver to assist 15-20 seniors at more affordable costs. By developing the AI optimization models to automate the complex and dynamic scheduling processes, the neighborhood model can be scaled repeatedly to help more seniors and facilitate mobile connectivity for the entire caregiving team, including family and professionals.

DATA

Analyze a sampling of de-identified data about routes and tasks for a hypothetical neighborhood of seniors. Develop optimization models to increase the efficiency of care by minimizing travel and downtime.

TECH APPROACH:

Develop mixed-integer optimization models, test and validate the optimized model vs. the hypothetical model and identify the magnitude of improvement and reduction of cost. Embed the automation into the digitized process of scheduling and dynamic capacity management to make operating a neighborhood consistently replicable.

PI(s): Dew-Anne Langcaon, CEO Vivia Cares

Kimia Ghobadi, PhD Johns Hopkins









Al for predicting adverse health events in the elderly population using wearable devices



AREA OF NEED:

Limited ability to continuously monitor and predict adverse health events in elderly patients, particularly in resource-constrained acute care settings, leading to delayed interventions and suboptimal outcomes.

OPPORTUNITY:

Develop and validate Al-driven monitoring solutions that can continuously assess patient status and predict adverse events, enabling early intervention and improved care management for elderly patients.

DATA:

Leveraging our SUMMIT data engine to process real wearable device data and simulated wearable data and develop machine learning models for real-time health monitoring and prediction, with focus on multimodal integration of various data types (waveforms, tables, clinical notes).


TECH APPROACH:

Working with MIMIC III waveform dataset, simulated data, and planning to collect additional data at two pilot sites, focusing on continuous physiological signals and clinical events. Using simulated data to augment data during training and evaluate foundation models for adaptation to physiological time series.


PI(s): Matthias Christenson, PhD

COHORT: GY3

MTN SUMMIT STACK



TRAINING APPROACH





Characterizing and Stratifying Cognitive Impairment Using Cognitive and Speech Al



AREA OF NEED: Diagnostic approaches for characterizing

AD/ADRDs rely on dichotomous "mild" versus "major" neurocognitive disorder categories, limiting the ability to capture subtle individual differences and develop more nuanced stratification of AD/ADRDs.

Develop precise, scalable methods to **OPPORTUNITY:**

capture baseline cognitive abilities using "gold standard" neuropsychological tests and remote digital tools to capture fluctuations in cognitive status.

Collect baseline neuropsychological data DATA:

at a single timepoint, along with cognitive

and speech EMA, and actigraphy monitoring 6 times over 1 month.

Combine gold standard **TECH APPROACH:**

neuropsychological testing with real-time digital assessments to capture within-

and between-person variability,

leveraging AI/ML/NLP algorithms (IBM


Digital Health) to refine diagnostic accuracy and support clinical decision-

making. Shifali Singh, Ph.D. PI(s):

COHORT:

Cognitive EMA

Using TestMyBrain



house

tree

Verbal Paired











Please click the microphone button and then say:

"The quick brown fox jumps over the lazy dog."

You may press the Stop button if you finish before the timer runs out.





Smartphone-Based Fall Prevention Therapy and the Israel Lahey Health Smartphone Based Fall Prevention Therapy and the Israel Deaconess Monitoring for Older Adults Medical Center

🖔 brightway

AREA OF NEED: Falls are among the greatest threats to

healthy aging. Fall-prevention physical therapy programs may be effective, but suffer from limits in accessibility, scalability,

and personalization.

OPPORTUNITY: Develop, validate, and test automated

delivery of balance assessments and fall prevention exercises using the Brightway

PT mobile application.

DATA: Collecting movement data for balance

assessments and fall prevention

exercises, as well stakeholder input and

user testing.

TECH APPROACH: Leverage computer vision models on the

Brightway PT platform, validate with laboratory-grade motion analysis, and prototype personalized real-time feedback for automated support to

users.

PI(s): Dennis Anderson, PhD (BIDMC)

Yannick Coehn (Brightway Health)





YayaGuide: Al-Enabled Personalized Training for Caregivers of Elders with ADRD

AREA OF NEED:

The demand both for skilled caregivers and for upskilling family caregivers is at an all-time high

OPPORTUNITY:

TECH APPROACH:

The mission of YayaGuide by CareYaya is to expand access to quality care by empowering caregivers of those living with

dementia.

DATA:

YayaGuide addresses this critical unmet need by harnessing the power of AI and micro-learning to provide personalized, engaging training at caregivers' fingertips – essentially, creating the "Duolingo of dementia care training". Coupled with conversational agents and machine learning for personalization, micro-learning can make

learning for personalization, micro-learning can make training highly accessible for time-strapped caregivers. It also promotes knowledge retention versus traditional

learning formats.

PI(s): Neal Shah





Frailty subtyping to identify paths to healthy aging


Area of Need: Everyone experiences aging uniquely, however we model frailty uniformally. A tailored approach to modeling frailty will facilitate the identification of personalized interventions to promote healthy aging.

Opportunity: In collaboration with the Rush Alzheimer's Disease Center, we have access to longitudinal data on over 5,000 individuals capturing aspects of their physical, cognitive, and social wellbeing over time.

Approach: We will develop an Al model to identify and predict frailty subtypes and characterize them macroscopically and molecularly. Clinicians may use our model to guide people toward a healthier state.

PI: Dr. Rebecca Keener

Cohort: GY4





In-Home Assessment of Muscle Deterioration with Al-Enhanced Smartphone Sonometry

AREA OF NEED:

Sarcopenia affects 6-22% of older adults, leading to muscle loss, falls, and hospitalizations. Current assessments are infrequent, require clinical visits, and miss early signs of deterioration.

OPPORTUNITY:

MusTrack enables in-home, frequent, and cost-effective muscle assessments. It allows early detection, longitudinal monitoring, and reduces healthcare costs by eliminating the need for specialized equipment.

DATA:


Collecting ultrasound B-mode imaging and smartphone ultrasound sonometry for muscle assessments


TECH APPROACH:

Al-enhanced smartphone sonometry uses metasurfaces and 3D printed cases for ultrasound sonometry. CycleGAN updates outdated B-mode imaging, while personalized models adapt handgrip strength tests to different users and devices.

PI(s):

Renjie Zhao, PhD (JHU) Xinyu Zhang, PhD (UCSD)







Expanding Epigenetic Analysis of Aging and AD/ADRD through Machine Learning

AREA OF NEED: Due to technical challenges, histone

modifications are typically not

experimentally profiled or considered in

cohort-based studies of aging and

AD/ADRD

OPPORTUNITY: There is the opportunity to

computationally impute histone

modification maps based on cohort DNA methylation array data and reference epigenomic data to study the biology of

aging and AD/ADRD

DATA: We will leverage reference epigenomic


data from the International Human Epigenome Consortium and cohort methylation array data focused on aging

and AD/ADRD

TECH APPROACH: We will investigate multiple machine

learning strategies to impute histone modification maps from array DNA methylation. We will conduct histone wide association study and epigenetic clock analyses based on the imputed data

PI(s): Jason Ernst, PhD (UCLA)





Feasibility and Acceptability of a Robotic Assistant for Early-Stage ADRD Care



AREA OF NEED:

The aging population and increasing prevalence of Alzheimer's Disease and Related Dementias (ADRD) pose challenges for caregiving, as staffing shortages and costs limit access to consistent, high-quality support. Robotic assistants may help by enhancing safety, independence, and social engagement for older adults living alone, whether in private residences or within assisted-living settings.

PROJECT GOAL:

Evaluate feasibility and acceptability of a prototype robotic aid from NaviGAIT Inc. for individuals with early-stage ADRD through real-world deployments, stakeholder input, and mixed-methods analysis.

METHODS:

Gathering qualitative and quantitative data through focus groups, supervised short visits, and extended overnight placements of the robotic assistant in residents' homes, along with insights from family caregivers and professional staff.


IMPACT:

Assessing robotic capabilities in real-world environments, including monitoring, reminders, and social engagement. Developing an evaluation framework to guide the design of robotic technologies that support aging in place.

INVESTIGATORS:

Philip A. Cola, Peter Whitehouse, and

Stefan Agamanolis





An Automated Non-Invasive Diagnostic Tool in Hydrocephalus Patients

AREA OF NEED: One of the most common problems in neurosurgery is

hydrocephalus: an accumulation of cerebrospinal fluid (CSF) causing elevated intracranial pressure. This occurs in many different forms but remains poorly understood. Tools are needed for improved, non-invasive diagnosis and guidance during management.

OPPORTUNITY: Develop and validate an automated deep learning-based approach

for ventricular and subarachnoid space segmentation with parcellation and quantitation applicable to NPH and non-NPH

patients over the age of 60.

DATA: Acquire retrospective multi-contrast magnetic resonance imaging

data from an internal cohort of ~800 NPH and suspected NPH

patients.

TECH APPROACH: Optimize and train VParNet, a previously developed deep network


for ventricular segmentation. Improve and optimize our


subarachnoid space segmentation method using deep networks, conventional deformable models, neural ordinary differential

equations, and physics-inspired neural networks.

PI(s): Jerry Prince, PhD (JHU)

Mark Luciano, MD, PhD (JHMI)







Al-Driven Cognitive Enhancement and Assessment Program for Adults at Risk for Alzheimer's Disease



AREA OF NEED: The increasing prevalence of Alzheimer's disease and other

dementias is a growing concern. While pharmacologic treatments are emerging, there remains a need for engaging, accessible, and

low-risk interventions that can be used at home.

OPPORTUNITY: Validate and test LookAlikes, a web-based cognitive enhancement

game where participants describe an image and receive immediate feedback to promote cognitive functions like memory, flexibility,

and planning.

DATA: Collecting verbal image descriptions, user engagement metrics,

and survey feedback from older adults with subjective memory concerns, along with data to validate the accuracy between image

and verbal descriptions.

TECH APPROACH: Use speech-to-text and image generation AI tools (Google Cloud

API and DALL·E), combined with image similarity scoring using a VGG16 model, to offer real-time feedback and monitor cognitive

function over time.

Pls: Meghan Mattos, PhD, RN (UVA)

Serkan Sandikcioglu, MS (Calbium AI)





MRI-based body composition to assess biological age and frailty



AREA OF NEED: C

Chronologic age is widely used in medical decision making but is a crude measure of aging. Better measures of "biologic aging" can improve clinical decisions and help researchers study how individuals age.

OPPORTUNITY:

Develop and test whether a tool can accurately predict biologic age based on body composition from MR images.

DATA:


Secondary analysis of >80,000 individuals with body MR imaging including the UK Biobank, the German National Cohort and Mass General Brigham patients.

TECH APPROACH:

Develop 3D computer vision models to identify muscle and fat regions and subsequently predict biologic age. Validate whether this predicted age is 1) associated with aging-related disease, 2) can be used to better understand how aging affects body composition.


PI(s): Vineet K Raghu, PhD

COHORT: GY4



Segmentation model identifies fat and muscle from body MR images

MRI-Age model estimates biologic age from segmented muscle and fat regions





Safe Circle:



Al and Micro-radar-based Remote Monitoring for Patients with AD/ADRD


AREA OF NEED:

SafeCircle addresses the urgent need for privacy-focused, accurate, and affordable remote monitoring for patients with AD/ADRD, tackling high risks of falls and wandering that strain caregivers and healthcare systems.



OPPORTUNITY:

The SafeCircle project aims to develop and evaluate an AI-enabled, micro-radar-based iOS prototype for accurately detecting falls and wandering in AD/ADRD patients, enhancing patient safety and reducing caregiver burden.



DATA:

Leverage non-identifiable micro-radar data capturing spatiotemporal movement patterns and event-triggered video snippets.

Data collection follows IRB-approved protocols.

PI(s):

Dr. Nazmus Sakib (KSU) Dr. Sumit Chakravarty (KSU)

COHORT:

GY4

TECH APPROACH:

It integrates FMCW micro-radar technology with deep learning models for accurate fall and wandering behavior detection, leveraging an Agile Scrum framework for iterative iOS prototype development.



Acceptability of Voice Clones in Agents for Aging Adult Wellness Monitoring



AREA OF NEED: Familiarity in caring for the elderly

population is of paramount

importance. The ability to provide consistent interaction results in improved outcomes, but time is

limited for most caregivers.

OPPORTUNITY: The goal of this study is to

understand how Virtual Agents,

including Voice Clones, can enhance digital interactions for older adults.

DATA: Collect survey data from participant

who experience interactions with voice clones of someone close to them with a reflection on their well

being

TECH APPROACH: Utilize generative AI technologies to

create voice clones of


caregivers/loved ones to study

impact on participants.

PI(s): Amit Mehta, MD

Camille Noufi, PhD

COHORT:





Patient-centered precision medicine lab result communication for older adults



AREA OF NEED: The status quo of lab result reporting fails

to provide older adults with personalized information that they can understand and use to make informed decisions about their

healthcare and health behaviors.

OPPORTUNITY: Create models to predict and

communicate the individualized 2- and 5-

year risk of developing ESKD for adults

aged 65 years and older.

DATA: The model will be trained on UC Health

patient data, with the implementation

guided by usability testing and

stakeholder engagement.


TECH APPROACH: Leverage machine learning models to

calculate risk from EHR records and communicate these personalized risks as

part of existing lab workflows and

processes.

PI(s): Catherine Sarkisian, MD, MSHS (UCLA)





Developing & Testing an Evidence-based, Al Dementia Care Navigation Assistant





AREA OF NEED: One of the best ways to improve outcomes

and reduce costs in dementia is with highquality care navigation to keep the patient at home. However, we have a shortage of

care navigators.

OPPORTUNITY: Develop, validate, and test a care

navigator support bot leveraging the Johns Hopkins MIND at Home program

and the Craniometrix chatbot.

DATA: Pre-created training data from the MIND

at Home program, and training feedback from care navigators leveraging the bot.

TECH APPROACH: Leverage Retrieval Augmented

Generation techniques to build a high-

quality chatbot to support care


navigators, preventing hallucinations by

requiring referential retrieval.

PI(s): Nikhil Patel (Craniometrix)

Halima Amjad, MD MPH PhD (Hopkins)

Cynthia Fields, MD (Hopkins)





Using Retitrack+AI to Explore Novel Eye Biomarkers in Patients with Cognitive Impairment



AREA OF NEED: Current AD/MCI diagnostics are invasive,

costly, and lack early sensitivity. There's a need for scalable, objective biomarkers.

OPPORTUNITY: Develop, validate, and test automated

detection of cognitive decline using highresolution retinal and pupillary eye movement data captured by the

Retitrack[™] device.

DATA: Retitrack[™] captures FEMs, saccades, and

pupillary dynamics. Cognivue Clarity provides cognitive performance metrics. All data is de-identified and securely

stored for AI analysis.

TECH APPROACH: Train a Transformer-based model on

synchronized eye movement data to

identify early signs of MCI/AD.

PI: Dr. Jon Artz (Renown Health)

COHORT: 104 adults with MCI or early dementia

from Renown Health Neurology Clinic **Inclusion**: Age 55-90, English-speaking,

VA ≥20/40, confirmed diagnosis

Exclusion: Other dementias, metabolic

issues, TBI







www.clighttechnologies.com

Machine learning to identify dermatoporosis in older adults

AREA OF NEED: Dermatoporosis, or skin frailty, affects over 30%

of adults age 65 years old or older and leads to easy bruising and skin tearing. It is associated other conditions such as bone health or skin cancer. However, older adults may not recognize their dermatoporosis and need for in-person care

OPPORTUNITY: Develop, validate, and test automated

assessments for dermatoporosis

DATA: Collect digital images from smartphones for

dermatoporosis assessments, as well stakeholder input and user testing.

TECH APPROACH: Leverage machine learning technologies to train,

validate and test skin images that have been labeled by dermatologists with skin frailty scores. Identify path forward so this tech can be accessible and empower older adults with skin frailty to seek in-person healthcare.

PI(s): Anne Lynn S. Chang MD Professor of Dermatology

Stanford University School of Medicine







JOHNS HOPKINS A Personalized Evolvable Al Tool for Real-Time **Testing and Intervening AD/ADRD**





AREA OF NEED:

Alzheimer's Disease (AD) and dementia resulting from it affect millions of people. Early diagnosis of AD is important but challenging. It can also be expensive, not always accessible, and costly.

OPPORTUNITY:

Develop, validate and test reliable selfadministered personalized assessments of the progression of AD based on easily available data through a mobile application.

DATA:

Relatively cheaper data like cognitive tests based on activities of daily living, speech, and optionally more expensive data such as MRI images.


TECH APPROACH:

Pre-trained AI models equipped with learning onthe-go to personalize the assessment. The cognitive tests are derived from everyday activities. Medical images and lab results are optionally used to enrich the prediction. Ensemble methods combine the predictions to improve performance.

PI(s):

Chung-Yi Chiu, PhD, ChengXiang Zhai, PhD (UIUC),

Yogatheesan Varatharajah, PhD (UMN)







Wearable in-shoe Sensors for Gait Monitoring in Normal Pressure Hydrocephalus (NPH)

- University of California, Irvine
- Johns Hopkins
- University of Florida

AREA OF NEED: Falls are among the greatest threats to healthy aging.

This is particularly problematic in patients with NPH where gait instability is a cardinal feature. Early detection can lead to early neurosurgical treatment

that can be transformative.

OPPORTUNITY: Develop, validate, and test in shoe sensors to find

pressure and gait patterns specific for NPH at its early stages. The shoe sensors may be used to track

improvement after surgery.

DATA: Collecting data from in-shoe sensors to characterize

the walk of NPH patients. We are looking for specific pressure point data and stride/pace characteristics that may lead to early identification of NPH patients.


TECH APPROACH: Leverage Artificial Intelligence/Machine Learning

models to analyze the large number of data points for gait derived from the in-shoe sensors. Analysis of changes in gait disorders over time with or without surgery will be used to develop a predictive model of


gait dysfunction in NPH.


PI(s): Jefferson Chen, MD, PhD (UCI)

Vadim Zipunnikov, PhD (JHU) Nikolay Bliznyuk, PhD (UF)











Identifying Digital Biomarkers for Early Cognitive Impairment



AREA OF NEED:

With aging of the population, the prevalence of cognitive impairments will increase. There's a need for accessible and scalable screening methods for detection of mild cognitive impairment (MCI) for earlier introduction of targeted pharmacologic and lifestyle interventions

OPPORTUNITY:

Develop, validate, and test machine learning models for detection of MCI risk in clinics and community settings

DATA:

Collecting data on eye movement and speech and comprehensive cognitive assessment, as well as user experience about data collection, from 150 older adults (75 MCI, 75 control)

TECH APPROACH:


Applying machine learning to data collected using iPad- and Tobii Glasses Pro, to develop sensitive algorithms for detection of MCI

PI(s):

Anis Davoudi, PhD (JHU)

COHORT: G'

GY5





Leveraging EHR and AI for High-Resolution ADRD Prevalence and Risk



AREA OF NEED:

Current Alzheimer's disease and related dementias (ADRD) surveillance relies on lagging, low-resolution estimates. This prevents timely identification of at-risk populations, is fragmented across systems, and cannot be scaled across states. Through the Federal BOLD program, states are attempting to develop basic ADRD surveillance systems, and Apriqot's approach helps them achieve this

goal.

OPPORTUNITY:

A next-generation surveillance model that integrates clinical and community data into high-resolution, near real-time estimates of ADRD prevalence and risk. This system will provide actionable insights for healthcare systems, state public health agencies, and community organizations, enabling earlier detection, better resource allocation, improved outcomes, and a replicable model for national use.

DATA:

Retrospective EHR data (2019–2024) from MaineHealth; de-identified population-level analyses; cohort: adults ≥65 with ≥2 visits (2020–2024), excluding prior dementia diagnosis/medication use.

TECH APPROACH:

Application of advanced AI and geospatial modeling to EHR data, integrated with demographic datasets. Development of predictive models that are portable across health systems, generating actionable dashboards and decision-support tools.

PI(s): Kevin Konty, PhD

COHORT: GY5

Population-level ADRD Risk





CAPTASK: Vision-Guided, Context-Aware Task Assistance for Daily Activities





AREA OF NEED: Alzheimer's Disease and Related Dementias

(AD/ADRD) significantly impact an individual's ability

to perform Activities of Daily Living (ADLs),

diminishing independence and quality of life while

placing immense burden on caregivers

OPPORTUNITY: Develop and evaluate an adaptable perceptual

artificial intelligence (AI) system that leverages vision-guided input to provide personalized task

assistance for individuals with AD/ADRD.

DATA: Collecting pilot data for task guidance in daily tasks

such as meal preparation and medication management.

TECH APPROACH: Leverage advances in large foundation models and

reasoning mechanisms to develop a proactive conversational interface that can provide

personalized assistance.

PI: Joyce Chai (Computer Science and Engineering, UM)

Anson Kairys (Psychiatry, UM)





Early Detection of Alzheimer's Using Non-Invasive Wearable Sensing and Al





AREA OF NEED: Alzheimer's Disease affects over 7 million

Americans, devastating lives with costs to exceed \$3.3 trillion/year by 2060. Current diagnostics are expensive, invasive and often inaccessible creating detection delays

and treatment monitoring difficulties.

OPPORTUNITY: Develop, validate, and test CogniTrend, a

cognitive health app that leverages wearable device data + AI to assess changes in cognitive wellness.

DATA: Collecting physiological and behavioral

data using wearable sensors from athome daily life behavior combined with standardized cognitive testing results.

TECH APPROACH: Leverage the latest advances in Al


combined with high-fidelity wearable sensing data to develop machine learning models that can predict cognitive

wellness.

PI: Sean Montgomery, PhD

(Connected Future Labs)









Integrating AI into a Smart Sleep Mask to Improve Slow Wave Sleep in AD Patients

AREA OF NEED:

Sleep disturbances are a common facet of Alzheimer's disease (AD) affecting 20-40% of patients in the mild to moderate stage of the illness. There is strong evidence that poor slow wave sleep accelerates the progress of AD through the failure to clear brain toxins during sleep.

OPPORTUNITY:

To reduce the accelerating impacts of disturbed sleep and slow the disease progression in AD patients, thereby extending quality of life and reducing the burden on caregivers.

DATA:

Collect EEG brain wave data for sleep classification, stimulate with dynamic peripheral neuromodulation patterns (TRS), and assess the improvement of slow wave sleep in a closed loop system.

TECH APPROACH:


Develop a comfortable sleep wearable medical device that monitors an AD patient's sleep and provides TRS non-invasive artificial intelligence-driven personalized peripheral neuromodulation to promote healthy slow wave sleep.

PI(s): Brian Krohn, PhD (Synaptic Health)

COHORT: GY5



Al Driven Sensing & Stimulating Feedback Loop







Smart Robot Guide for Daily Care in People with Memory Loss



AREA OF NEED:

People with memory loss struggle with hydration, oral hygiene, and nutrition—tasks making up most daily caregiving—yet current technologies fail to sustain engagement as cognitive decline progresses.

OPPORTUNITY:

Develop and pilot an Al-powered humanoid robot to deliver personalized, real-time multimodal support for hydration, oral hygiene, and nutrition in people living with dementia.


DATA:

Collecting video and sensor data on user-robot interactions in daily living spaces, alongside user testing and stakeholder feedback.

TECH APPROACH:

Integrate Al-powered Pepper robot with unobtrusive sensing and multimodal perception (LLMs, Affective computing) for adaptive ADL support in dementia care through embodied, context-aware assistance.

PI(s): Fiona Yuan, PhD, Worcester Polytechnic Institute





Re-Kinesis: Wearable Gait Analysis Laboratory for Aging Populations



AREA OF NEED:

Gait impairment is common in stroke and aging. Clinics rely on eyeballing, while gait labs—though comprehensive—are costly and hard to access. Patients lack feedback, and clinicians lack continuous insight. A device is needed to deliver accessible 2D plantar and 3D motion data with remote monitoring and actionable insights to track progress and guide care.

OPPORTUNITY:

Develop and validate *Re-Kinesis*, a lightweight in-shoe sensing system that transforms everyday walking into objective mobility data. The system translates 2D plantar dynamics into 3D lower limb motion, bringing gait lab-quality analysis as a new vital sign into routine rehab visits and daily living.

DATA:


High-resolution plantar pressure recordings combined with Aldriven kinematic reconstruction. Clinical validation with older adults in the AITC registry across two study visits, paired with feedback from rehabilitation physicians and patients.

TECH APPROACH:

Optimize and stress-test wearable hardware and user interface for users, validate accuracy against gold-standard gait lab measures, and refine usability for real-world rehabilitation settings. Package results and workflow documentation to support future SBIR/STTR commercialization.

PI: Nitish V. Thakor, PhD (Johns Hopkins University)







A Morphologic-Epigenomic Tool Mapping Senotherapy Responses to Senescent Cells AtlasXomics

Cellular senescence is implicated for both **AREA OF NEED:**

healthy aging and age-related diseases. Yet

methods to monitor and identify

senescence subtypes ex vivo and within

tissue are lacking.

Develop, validate, and test a combined, spatial **OPPORTUNITY:**

morphological and epigenomic assay for the purposes of, 1) screening new therapies

affecting senescence subtypes, 2) identifying

senescence subtypes in tissues, and 3) identify

effective senotherapies per tissue

Collect and correlate senescence-DATA:

induced cell morphologies and

epigenomic data from spatial tissue at

the single-cell level.

Leverage Machine Learning to classify **TECH APPROACH:**

> functional subtypes driving senescent phenotypes, identifying lead epigenomic pathways to target for novel therapies in


age-related diseases and ultimately

ADRD.

PI(s): Timothy McConnell (AtlasXomics)

Jude Phillip (JHU)

COHORT: GY5




Jude Phillip Lab


Established Senescence Subtyping Model

Aim 2






Single-cell Epigenomics



Aim 2

Tissue Epigenomics



AtlasXomics Commercial Spatial **Epigenomics Platform**



Al-based mitochondrial multimodal aging clock and biomarkers





AREA OF NEED:

Epigenetic clocks based on nuclear DNA are among the best molecular methods to estimate biological age. However, the accuracy of estimating chronological age degrades at older ages and few clocks have strong predictive power for health status. Mutations in mitochondrial DNA (mtDNA) contribute to their mitochondrial dysfunction, considered a hallmark of aging. In contrast to mutations, little is known about the epigenetic regulation of mtDNA. Little work has been done to develop aging clocks based on mtDNA.

OPPORTUNITY:

We propose to develop AI-based mitochondrial aging clock models using mtDNA mutational and epigenomic data.

DATA:

Sequence mitochondrial DNA (mtDNA) samples from approximately 80 individuals, ranging in age from 20 to >89 years (10 samples per decade age group) using Oxford Nanopore Technologies long-read sequencing to capture DNA mutations and methylation (5mC, 5hmC, and 6mA).

TECH APPROACH:

Leverage linear and nonlinear Al algorithms to integrate genomic and epigenomic mutations and develop a highly accurate mitochondrial aging clock.

PI(s): Hayan Lee, PhD (Fox Chase Cancer Center)

Glenn Gerhard, MD (Lewis Katz School of Medicine, Temple Univ)

